

Medivit

Vitamin B-Complex

COMPOSITION:

Medivit Tablet: Each film coated tablet contains Thiamine Mononitrate BP 5 mg, Riboflavin BP 2 mg, Pyridoxine Hydrochloride BP 2 mg and Nicotinamide BP 20 mg.

PHARMACOLOGY:

Medivit is a balanced combination of important members of vitamin B complex. It contains vitamins that are chemical compounds, which are essential for health. They act primarily as regulators of metabolic processes and play a role in energy transformations, usually acting as co-enzymes in enzymatic systems.

INDICATION:

Vitamin B-complex deficiency states as-

- a. Nutritional deficiency anaemias
- b. During pregnancy and lactation
- c. Pre- and post-operative conditions
- d. During antibiotic therapy
- e. Chronic constipation, stomatitis
- f. Liver disease
- g. Convalescence
- h. Hyperemesis gravidarum and patients on restricted diets.

DOSAGE AND ADMINISTRATION:

Medivit Tablet: 1-2 tablets 3 times daily, or as recommended by the registered physician.

CONTRAINDICATION:

Medivit should not be taken in case of hypervitaminosis and hypersensitivity to any of its ingredients.

WARNING & PRECAUTIONS:

Caution should be taken during concomitant use of vitamin B-complex and levodopa.

SIDE EFFECT:

No significant side effects have been reported.

USE IN PREGNANCY AND LACTATION:

Safety margin has been established in taking **Medivit** during pregnancy and lactation.

OVERDOSAGE:

Excess vitamin B is readily excreted; therefore no serious problems are anticipated for the administration of vitamin B in this form.

STORAGE:

Store in a cool and dry place (below 30 °C), protected from light and moisture. Keep out of reach of children.

PACKING:

Medivit Tablet: Each box contains 10x10 tablets in blister pack.

Manufactured by:



MEDICON Pharmaceuticals Ltd.
Mirpur, Dhaka, Bangladesh